

### Comparison between HIRDLS and MIPAS



#### Radiances and Key Retrieved Species



Claire Waymark

University of Oxford: Anu Dudhia, John Barnett and Fred Taylor University of Colorado: John Gille, Cheryl Craig and Rashid Khosravi

+ HIRLDS team


+ MIPAS team


EOS AURA


Chemistry & Climato

## The Michelson Interferometer for Passive Atmospheric Sounding (MIPAS)

- Launched on ESA's ENVISAT in March 2002
- Fourier Transform spectrometer
- Measures in the infrared
- Uses limb geometry
- Scans sequentially through atmosphere
- Routine products are pressure temperature, O<sub>3</sub>, H<sub>2</sub>O, HNO<sub>3</sub>, CH<sub>4</sub>, N<sub>2</sub>O and NO<sub>2</sub>.
- Reduced resolution of 0.0625 cm<sup>-1</sup>








# The High-Resolution Dynamic Limb Sounder (HIRDLS)





- Launched on AURA in July 2004
- Limb geometry
- Measures in the infrared
- Radiometer fixed spectral channels
- Each channel targets a specific species
- Obstruction cover all but ~20% on one viewing angle.
  - Planed routine products are pressure temperature, 03, H2O, HNO3, CH4, N2O, NO2, N2O5, ClONO2, F11 and F12








#### HIRDLS and MIPAS



MIPSAS and HIRDLS L1B CSepage for the 44 pos




♦ HIRDLS day

Aura Science Meeting, Boulder 11-15 Sep 06



### HIRDLS and MIPAS Spectral Coincidences





- 10 HIRDLS channels completely overlap the MIPAS spectrum.
- No temperature channel overlap.





#### Radiance Comparison: Channels 8





Residuals (MIPAS - HIRDLS )/MIPAS

28<sup>th</sup> January 2005









- Comparisons use radiances that go into the v2 processing
- Except for dashed residuals which use radiances used for a previous internal processing version
- Different colour residuals represent the latitude bands 20-60N, 205-20N and 20-60S.



20-60N 205-20N 20-60S

HIRDLS.



ting, Boulder 11-15 Sep 06




#### Radiance Comparison: Channel 18



Residuals (MIPAS - HIRDLS )/MIPAS


28<sup>th</sup> January 2005




4th May

2006

(20N-60N) HIRDLS Channel 18 (H20)





- Comparisons use radiances that go into the v2 processing.
- Except for dashed residuals which use radiances used for a previous internal processing version

Different colour residuals represent the latitude bands 20-60N, 205-20N and 20-60S.



0.0001 0.0010 0.0100 0.1000
Radiance (W/M-2/Sr)

HIRDLS.



ting, Boulder 11-15 Sep 06




#### Temperature Climatology comparison



HIRDLS



MIPAS Climatology





Science Meen



### Temperature retrievals










### Ozone retrieval comparison








### In-direct radiance comparison







### MIPAS / HIRDLS Altitude Shift



- Least squares fit method used
- Calculations using the 28<sup>th</sup> January data set
- Altitude shift calculated from direct comparisons in black
- Altitude shift calculated from indirect comparisons in aqua









- ullet There is good agreement between the HIRDLS and MIPAS zonally averaged radiances .
- Seems to be a gain offset for the 4<sup>th</sup> May data.
- The Temperature and O<sub>3</sub> retrievals agree well for 28<sup>th</sup> January 2005 and 4<sup>th</sup> May 2006. More detailed structure seen for 4<sup>th</sup> May 06
- In-direct radiance comparisons show good agreement in for the temperature channels

#### Future work

- Calculate an altitude shift for 4<sup>th</sup> may 06.
- Run the comparisons using v2.02.

## Thank you for your attention